

Hints of Integrability Beyond the Planar Limit: Nontrivial Backgrounds

Robert de Mello Koch^{1,2} Tanay K. Dey¹, Norman Ives¹ and Michael Stephanou¹

¹ *National Institute for Theoretical Physics,
Department of Physics and Centre for Theoretical Physics,
University of the Witwatersrand,
Wits, 2050,
South Africa*

² *Stellenbosch Institute for Advanced Studies,
Stellenbosch,
South Africa*

E-mail: robert@neo.phys.wits.ac.za, Tanay.Dey@wits.ac.za,
Norman.Ives@students.wits.ac.za,
Michael.Stephanou@students.wits.ac.za

ABSTRACT: The problem of computing the anomalous dimensions of a class of (nearly) half-BPS operators with a large $\mathcal{R}$ -charge is reduced to the problem of diagonalizing a Cuntz oscillator chain. Due to the large dimension of the operators we consider, non-planar corrections must be summed to correctly construct the Cuntz oscillator dynamics. These non-planar corrections do not represent quantum corrections in the dual gravitational theory, but rather, they account for the backreaction from the heavy operator whose dimension we study. Non-planar corrections accounting for quantum corrections seem to spoil integrability, in general. It is interesting to ask if non-planar corrections that account for the backreaction also spoil integrability. We find a limit in which our Cuntz chain continues to admit extra conserved charges suggesting that integrability might survive.

KEYWORDS: AdS/CFT correspondence, super Yang-Mills theory.

Contents

1. Introduction	1
2. Two Loop Cuntz Chain of the LLM Background	3
2.1 Annulus Background	3
2.2 Two Loop Effective Dilatation Operator	4
2.3 Action of the Two Loop Effective Dilatation Operator	6
2.4 Leading $M + N$ Limit	8
3. Conservation Laws	11
3.1 U_2 rewritten in the Cuntz Oscillator Language	12
3.2 Classical Limit	13
3.3 Integrability in the Large M Limit	17
4. Discussion	19
A. Two Point Functions	19
B. More on the Cuntz Chain	21
C. Explicit Expressions for the Two Loop Dilatation Operator	22

1. Introduction

There is by now an impressive body of work suggesting that *planar* $\mathcal{N} = 4$ super Yang-Mills theory is *exactly integrable*. This would be very fortunate indeed, since it would mean the problem of computing the spectrum of all possible scaling dimensions of the gauge theory can be solved exactly, in the large N limit, by employing a Bethe ansatz. This has been established for the complete set of possible operators at one loop in [1, 2] and to two and three loops in the $\text{su}(2)$ sector [3]. Given these results it is natural to guess that integrability extends to all orders in perturbation theory and perhaps even to the non-perturbative level [3]. There is now mounting evidence that this guess is correct [4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16].

The idea of spin chain parity played a central role in the discovery of the planar two and three loop integrability[3]. Acting on a single trace operator, parity simply reverses the order of fields inside the trace. The dilatation operator commutes with parity, so that as we would expect, the dilatation eigenstates are also parity eigenstates. In addition, [3] found that eigenstates with opposite parity were degenerate - this was quite unexpected. This degeneracy can be explained by the existence of a higher conserved charge (U_2 in the notation of [3]) that commutes with the dilatation operator but anticommutes with parity. Its also possible (and useful) to put this argument on its head: one can interpret the degeneracy of states with opposite parity as evidence for the existence of further conserved quantities. When non-planar corrections are taken into account, parity remains a good quantum number but the degeneracy is lifted (see [3] for a discussion of the $\mathcal{N} = 4$ super Yang-Mills case and [17] for a very relevant discussion in the context of the ABJM theory). Although this only proves that the standard construction of conserved charges does not work away from the strict planar limit, it does suggests that integrability might not survive away from this limit. Clearly an important question is

Does integrability survive non-planar corrections?

In this article we will explicitly describe a situation in which we do sum (in fact, an infinite number of) non-planar corrections. Further, we collect some evidence that the resulting dynamics remains integrable. This suggests that, at least in certain situations, the answer to the above question is positive.

The non-planar corrections that we consider arise because we are interested in computing the anomalous dimensions of operators whose classical dimension is of order N^2 . The usual $1/N$ suppression of non-planar diagrams is, in this case, overpowered by huge combinatoric factors[18]. In a series of articles[19, 21, 22], building on the earlier works[23, 24, 25], we have developed techniques to systematically study these operators. The specific operators we study are spelt out in detail in section 2.1. In section 2.2 we give the dilatation operator to two loops. To obtain this result, ribbon diagrams with arbitrarily large genus are summed. In section 2.3 we study the action of our dilatation operator. The action of this dilatation operator is *not easily formulated in a spin chain language because, nonplanar corrections allow the number of fields within each trace to change*. This translates into a spin chain with a variable lattice length. A convenient reformulation of the problem, in terms of a Cuntz lattice, was given in [26]. In the reformulation, particles (described by Cuntz oscillators) hop on a lattice of fixed size. The fact that the size of the spin chain lattice was dynamical now translates into the fact that the total number of particles populating the Cuntz lattice is dynamical. We give the Cuntz oscillator description of the dilatation operator for the class of operators

we study in section 2.4. In section 3 we start to look for signs that our Cuntz chain is integrable. To start, we rewrite the spin chain (in the $\mathfrak{su}(2)$ sector) of [1, 2, 3] in the Cuntz oscillator language. In particular, we write the conserved charge U_2 of [3] in terms of Cuntz oscillators. We have verified, using this expression for U_2 , that U_2 does not commute with the Cuntz chain Hamiltonian corresponding to the annulus geometry. Another way to explore the integrability of the original model is to study the semi-classical limit of the spin chain, which can be matched to the low energy limit of the principal chiral model. It is known that the principal chiral model is integrable. We can show that the Cuntz chain corresponding to the spin chain of [1, 2, 3] is indeed equivalent to the low energy limit of the principal chiral model - the spin chain and the Cuntz chain simply correspond to different choices of gauge. We give the explicit form of the gauge transformation relating the two in section 3.2. In section 3.3 we study the large M limit of our Hamiltonian and argue that we can indeed write down higher conserved charges. This suggests that integrability survives in this limit. In section 4 we discuss our results.

2. Two Loop Cuntz Chain of the LLM Background

$\mathcal{N} = 4$ super Yang-Mills theory has 6 scalars ϕ_i transforming in the adjoint of the gauge group and in the **6** of the $SU(4)_{\mathcal{R}}$ symmetry. We shall use the complex combinations

$$Z = \phi_1 + i\phi_2, \quad Y = \phi_3 + i\phi_4, \quad X = \phi_5 + i\phi_6, \quad (2.1)$$

in what follows. All operators that we study are built using only Z and Y ; they belong to the $\mathfrak{su}(2)$ sector of the theory. Many of the expressions that we write involve traces over Z s, Y s and derivatives of them. To avoid confusion, we will now spell out the index structure of a few expressions

$$\text{Tr} \left(Z \frac{\partial}{\partial Z} \right) = Z_{ij} \frac{\partial}{\partial Z_{ij}}, \quad (2.2)$$

$$\text{Tr} \left(ZY \frac{\partial}{\partial Z} \frac{\partial}{\partial Y} \right) = Z_{ij} Y_{jk} \frac{\partial}{\partial Z_{lk}} \frac{\partial}{\partial Y_{il}}. \quad (2.3)$$

2.1 Annulus Background

Schur polynomials provide a very convenient reorganization of the half-BPS sector of the $\mathcal{N} = 4$ super Yang-Mills theory. This is due to the fact that their two point function is known to all orders in $\frac{1}{N}$ [23, 24] and that they satisfy a product rule allowing

computation of exact n -point correlators using only two-point functions¹. The half-BPS sector of the theory can be reduced to the dynamics of the eigenvalues of Z , which is the dynamics of N non interacting fermions in an external harmonic oscillator potential[23, 28]. The half-BPS sector of operators with $\mathcal{R}$ charge of order N^2 , are dual to solutions of type IIB supergravity - the LLM geometries[29]. For a careful discussion of precisely what aspects of the eigenvalue dynamics the supergravity captures and vice versa, see [31]. For a recent discussion of the macroscopic description of the dual geometry see [32]. It is by now well known that the space of the LLM geometries is given by a black and white coloring of a two dimensional plane[29, 30]; this colored plane is isomorphic to the fermionic phase space of the eigenvalues of Z . The Schur polynomials correspond to geometries that are invariant under rotations in this plane.

The Schur polynomials are labeled by Young diagrams. In what follows, we will be interested in $\chi_B(Z)$ with B a Young diagram that has N rows and M columns. We take M to be of order N . Note that we can also express

$$\chi_B(Z) = (\det(Z))^M. \quad (2.4)$$

This implies that

$$\frac{\partial}{\partial Z_{ij}} \chi_B(Z) = M(Z^{-1})_{ji} \chi_B(Z), \quad (2.5)$$

a formula that we will make good use of below. The coloring describing the dual LLM geometry is a black annulus. The inner white disk has an area of $\frac{M\pi}{N}$ whilst the black annulus itself has an area of π in units that assign an area of $\frac{\pi}{N}$ to each fermion state in phase space.

2.2 Two Loop Effective Dilatation Operator

The two loop dilatation operator, in the $\mathfrak{su}(2)$ sector, has been computed in [3]. Using the conventions of [3], the dilation operator can be expanded as

$$D = \sum_{k=0}^{\infty} \left(\frac{g_{YM}^2}{16\pi^2} \right)^k D_{2k} = \sum_{k=0}^{\infty} g^{2k} D_{2k}, \quad (2.6)$$

where the tree level, one loop and two loop contributions are

$$D_0 = \text{Tr} \left(Z \frac{\partial}{\partial Z} \right) + \text{Tr} \left(Y \frac{\partial}{\partial Y} \right), \quad (2.7)$$

¹There are generalizations of these results to multimatrix models[27]. These results will be very relevant for studies of backgrounds that preserve less supersymmetry.

$$D_2 = -2 : \text{Tr} \left([Z, Y] \left[\frac{\partial}{\partial Z}, \frac{\partial}{\partial Y} \right] \right) :, \quad (2.8)$$

$$\begin{aligned} D_4 = & -2 : \text{Tr} \left(\left[[Y, Z], \frac{\partial}{\partial Z} \right] \left[\left[\frac{\partial}{\partial Y}, \frac{\partial}{\partial Z} \right], Z \right] \right) : \\ & -2 : \text{Tr} \left(\left[[Y, Z], \frac{\partial}{\partial Y} \right] \left[\left[\frac{\partial}{\partial Y}, \frac{\partial}{\partial Z} \right], Y \right] \right) : \\ & -2 : \text{Tr} \left([[Y, Z], T^a] \left[\left[\frac{\partial}{\partial Y}, \frac{\partial}{\partial Z} \right], T^a \right] \right) : . \end{aligned} \quad (2.9)$$

The normal ordering symbols here indicate that derivatives within the normal ordering symbols do not act on fields inside the normal ordering symbols.

We allow this dilatation operator to act on an operator built mainly from Z s with a few “impurities” = Y s added. For most of our study we explicitly display formulas for operators with two or three impurities. Our final formulas are however, completely general, covering the case that we have $O(1)$ impurities. Adapting the notation of [3] we define

$$\mathcal{O}_B(p, J_0, J_1, \dots, J_k) \equiv \chi_B(Z) \mathcal{O}_p^{J_0; J_1, \dots, J_k} \equiv \chi_B(Z) \text{Tr} (Y Z^p Y Z^{J_0-p}) \prod_{i=1}^k \text{Tr} Z^{J_i}, \quad (2.10)$$

$$\mathcal{Q}_B(J_0, J_1, J_2, \dots, J_k) \equiv \chi_B(Z) \mathcal{Q}^{J_0; J_1; J_2, \dots, J_k} \equiv \chi_B(Z) \text{Tr} (Y Z^{J_0}) \text{Tr} (Y Z^{J_1}) \prod_{i=2}^k \text{Tr} Z^{J_i}, \quad (2.11)$$

where $\chi_B(Z)$ is the operator creating the background, which was defined in the previous section. Our strategy is to define an effective dilatation operator D_{eff} as

$$D (\chi_B(Z) \mathcal{O}_p^{J_0; J_1, \dots, J_k}) = \chi_B(Z) D_{\text{eff}} \mathcal{O}_p^{J_0; J_1, \dots, J_k}. \quad (2.12)$$

Diagonalizing the action of D_{eff} on the gauge invariant operators $\mathcal{O}_p^{J_0; J_1, \dots, J_k}$ is clearly equivalent to diagonalizing the action of D on $\mathcal{O}_B(p, J_0, J_1, \dots, J_k)$. It is natural to interpret D_{eff} as the dilatation operator for the LLM background, that is, for the theory that is deformed by the insertion of $\chi_B(Z) \chi_B(Z^\dagger)$ in the path integral. Notice that we can write

$$D_{\text{eff}} = \frac{1}{\chi_B(Z)} D \chi_B(Z). \quad (2.13)$$

This formula remains correct even after replacing $\chi_B(Z)$ by any other operator creating the background, which depends only on Z . Ultimately, we will restrict ourselves to the large $M + N$ limit. To capture this limit, as explained in [22] one needs to resum an infinite number of nonplanar diagrams; ribbon diagrams with arbitrarily large genus

are summed. *This limit is certainly not the planar limit of $\mathcal{N} = 4$ super Yang Mills theory.*

The crucial observation needed in the computation of D_{eff} is that

$$\frac{\partial}{\partial Z_{ij}} (\chi_B(Z) \mathcal{O}_p^{J_0; J_1, \dots, J_k}) = \chi_B(Z) \left(\frac{\partial}{\partial Z_{ij}} + M(Z^{-1})_{ji} \right) \mathcal{O}_p^{J_0; J_1, \dots, J_k}. \quad (2.14)$$

Repeated application of this formula gives

$$D_{0 \text{ eff}} = D_0 + MN, \quad (2.15)$$

$$D_{2 \text{ eff}} = D_2 - 2M \text{Tr} \left((ZY Z^{-1} + Z^{-1} Y Z - 2Y) \frac{\partial}{\partial Y} \right), \quad (2.16)$$

$$\begin{aligned} D_{4 \text{ eff}} = & D_4 + 4NM \text{Tr} \left((ZY Z^{-1} + Z^{-1} Y Z - 2Y) \frac{\partial}{\partial Y} \right) \\ & + 2M : \text{Tr} [Z, Y] \left[Z^{-1}, \left[Z, \left[\frac{\partial}{\partial Z}, \frac{\partial}{\partial Y} \right] \right] \right] : + 2M : \text{Tr} [Z, Y] \left[\frac{\partial}{\partial Z}, \left[Z, \left[Z^{-1}, \frac{\partial}{\partial Y} \right] \right] \right] : \\ & + 2M : \text{Tr} [Z, Y] \left[\frac{\partial}{\partial Y}, \left[Y, \left[Z^{-1}, \frac{\partial}{\partial Y} \right] \right] \right] : \\ & - 2M^2 \text{Tr} \left((Z^2 Y Z^{-2} - 4ZY Z^{-1} + 6Y - 4Z^{-1} Y Z + Z^{-2} Y Z^2) \frac{\partial}{\partial Y} \right) \end{aligned} \quad (2.17)$$

for the tree level, one loop and two loop contributions to D_{eff} . The formula for $D_{0 \text{ eff}}$ has a straight forward interpretation - the dimension of the gauge invariant operator $\mathcal{O}_p^{J_0; J_1, \dots, J_k}$ is shifted by MN due to the presence of the background, which has dimension MN . The answer for $D_{2 \text{ eff}}$ has already been obtained and discussed in [33, 19, 21].

2.3 Action of the Two Loop Effective Dilatation Operator

A useful observation made in [3], is that, when acting with $g^2 D_2 + g^4 D_4$ on the generic two impurity gauge invariant operators $\mathcal{O}_B(p, J_0, J_1, \dots, J_k)$ and $\mathcal{Q}_B(J_0, J_1; J_2, \dots, J_k)$ operators of type $\mathcal{Q}_B(J_0, J_1; J_2, \dots, J_k)$ are never produced. This is easy to understand: acting with $g^2 D_2 + g^4 D_4$ always inserts a commutator $[Y, Z]$ into a trace; this trace vanishes unless it contains another Y . This observation generalizes: when acting with $g^2 D_{2 \text{ eff}} + g^4 D_{4 \text{ eff}}$ on the generic two impurity gauge invariant operators $\mathcal{O}_p^{J_0; J_1, \dots, J_k}$ and $\mathcal{Q}^{J_0, J_1; J_2, \dots, J_k}$ operators of type $\mathcal{Q}^{J_0, J_1; J_2, \dots, J_k}$ are never produced. This follows because both

$$\text{Tr} \left((ZY Z^{-1} + Z^{-1} Y Z - 2Y) \frac{\partial}{\partial Y} \right) \quad (2.18)$$

and

$$\text{Tr} \left((Z^2 Y Z^{-2} - 4 Z Y Z^{-1} + 6 Y - 4 Z^{-1} Y Z + Z^{-2} Y Z^2) \frac{\partial}{\partial Y} \right) \quad (2.19)$$

annihilate gauge invariant operators containing a single Y and because the remaining terms in $g^2 D_{2 \text{ eff}} + g^4 D_{4 \text{ eff}}$ always insert a $[Y, Z]$ into a trace. In what follows we will study the anomalous dimensions of the operators $\mathcal{O}_B(p, J_0, J_1, \dots, J_k)$. Clearly, from the observation we just made, we do not need to consider the operators $\mathcal{Q}_B(J_0, J_1, J_2, \dots, J_k)$ to do this.

Consider the action of $D_{2 \text{ eff}}$ on $\mathcal{O}_p^{J_0; J_1, \dots, J_k}$. Following [3] we can break $D_{2 \text{ eff}}$ into three pieces

$$D_{2 \text{ eff}} = D_{2,0 \text{ eff}} + D_{2,+ \text{ eff}} + D_{2,- \text{ eff}}, \quad (2.20)$$

where $D_{2,0 \text{ eff}}$ preserves the number of traces in $\mathcal{O}_p^{J_0; J_1, \dots, J_k}$, $D_{2,+ \text{ eff}}$ increases (by 1) the number of traces and $D_{2,- \text{ eff}}$ decreases (by 1) the number of traces. From (2.16) it is clear that the additional term proportional to M can only contribute to $D_{2,0 \text{ eff}}$. The terms $D_{2,+ \text{ eff}}$ and $D_{2,- \text{ eff}}$ which involve gauge invariant operator splitting and joining will not contribute at the leading order; they will be important when computing, for example, subleading corrections to the leading $M + N$ limit. The additional contributions proportional to M in (2.16) have an important effect: they imply that p and $J_0 - p$ can be negative. Thus, we need to consider gauge invariant operators in which we populate the two “gaps between the Y s” with both positive and negative powers of Z . This implies that p in $\mathcal{O}_p^{J_0; J_1, \dots, J_k}$ is completely unrestricted.

It is easy to write down an exact expression for the action of $D_{2 \text{ eff}}$

$$\begin{aligned} D_{2,0 \text{ eff}} \mathcal{O}_p^{J_0; J_1, \dots, J_k} = & -4A_1 \left(\mathcal{O}_{p+1}^{J_0; J_1, \dots, J_k} \right. \\ & \left. - \mathcal{O}_p^{J_0; J_1, \dots, J_k} \right) - 4A_2 \left(\mathcal{O}_{p-1}^{J_0; J_1, \dots, J_k} \right. \\ & \left. - \mathcal{O}_p^{J_0; J_1, \dots, J_k} \right), \end{aligned} \quad (2.21)$$

$$\begin{aligned} D_{2,+ \text{ eff}} \mathcal{O}_p^{J_0; J_1, \dots, J_k} = & 4 \sum_{J_{k+1}=1}^{p-1} \left(\mathcal{O}_{p-J_{k+1}}^{J_0-J_{k+1}; J_1, \dots, J_k, J_{k+1}} \right. \\ & \left. - \mathcal{O}_{p-J_{k+1}-1}^{J_0-J_{k+1}; J_1, \dots, J_k, J_{k+1}} \right) - 4 \sum_{J_{k+1}=1}^{J_0-p-1} \left(\mathcal{O}_{p+1}^{J_0-J_{k+1}; J_1, \dots, J_k, J_{k+1}} \right. \\ & \left. - \mathcal{O}_p^{J_0-J_{k+1}; J_1, \dots, J_k, J_{k+1}} \right), \end{aligned} \quad (2.22)$$

$$D_{2,- \text{ eff}} \mathcal{O}_p^{J_0; J_1, \dots, J_k} = 4 \sum_{i=1}^k J_i \left(\mathcal{O}_{p+J_i}^{J_0+J_i; J_1, \dots, \hat{J}_i, \dots, J_k} \right)$$

$$\begin{aligned}
& -\mathcal{O}_{p+J_i-1}^{J_0+J_i;J_1,\dots,\hat{J}_i,\dots,J_k} \Big) - 4 \sum_{i=1}^k J_i \Big(\mathcal{O}_{p+1}^{J_0+J_i;J_1,\dots,\hat{J}_i,\dots,J_k} \\
& \quad - \mathcal{O}_p^{J_0+J_i;J_1,\dots,\hat{J}_i,\dots,J_k} \Big), \tag{2.23}
\end{aligned}$$

where in the last expression hatted variables are removed from the argument of $\mathcal{O}$ and

$$\begin{aligned}
A_1 &= M + N & J_0 > p \\
&= M & \text{otherwise}
\end{aligned} \tag{2.24}$$

$$\begin{aligned}
A_2 &= M + N & p > 0 \\
&= M & \text{otherwise.}
\end{aligned} \tag{2.25}$$

Again motivated by [3] we can write

$$D_{4 \text{ eff}} = -\frac{1}{4}(D_{2 \text{ eff}})^2 + \delta D_{4 \text{ eff}} \tag{2.26}$$

where

$$\delta D_{4 \text{ eff}} = 2 : \text{Tr} [Z, Y] \left[\frac{d}{dY}, \left[Y, \left[\frac{d}{dZ}, \frac{d}{dY} \right] \right] \right] : + 2M : \text{Tr} [Z, Y] \left[\frac{d}{dY}, \left[Y, \left[Z^{-1}, \frac{d}{dY} \right] \right] \right] : .$$

We can decompose $\delta D_{4 \text{ eff}}$ as

$$\delta D_{4 \text{ eff}} = \delta D_{4,0 \text{ eff}} + \delta D_{4,+ \text{ eff}} + \delta D_{4,- \text{ eff}} + \delta D_{4,+- \text{ eff}} + \delta D_{4,-- \text{ eff}} + \delta D_{4,++ \text{ eff}} . \tag{2.27}$$

The number of pluses/minuses in the subscripts on the right hand side of this last expression tell us how many traces are added/removed by the action of that particular term. $\delta D_{4,+- \text{ eff}}$ adds and removes a trace and hence it comes from summing higher genus ribbon diagrams which contribute to the trace conserving piece of $\delta D_{4 \text{ eff}}$. It is again easy to write down exact expressions for the action of these terms. These expressions are given in Appendix C.

2.4 Leading $M + N$ Limit

To extract the leading terms at large N (recall that we take M to be of order N) we need to rewrite the action of $D_{2 \text{ eff}}$ obtained in the last subsection in terms of normalized gauge invariant operators, that is, gauge invariant operators that have a suitably normalized two point function. The relevant two point correlators have been computed in Appendix A. The result is most easily written in terms of a Cuntz oscillator chain. We will now focus on gauge invariant operators that have $k = 0$. If we relax the

restriction to two impurities (which we do from now on), the gauge invariant operators we study have the form

$$\mathcal{O}_B(\{p\}, J_0) \equiv \chi_B(Z) \text{Tr} (Y Z^{p_1} Y Z^{p_2} \dots Y Z^{p_n}), \quad \sum_{i=1}^n p_i = J_0. \quad (2.28)$$

To translate this gauge invariant operator into a Cuntz chain state, we associate a site of the Cuntz chain with each of the gaps between the Y s. The above gauge invariant operator defines a state in a Cuntz chain with n sites. Further, the number of Z s in each site gives the occupation number of that site. Finally, we require that operators with a normalized (free field) two point function map into normalized Cuntz chain states. This last point deserves a few comments. The spacetime dependence of the free field correlators we compute is trivially determined by the bare dimension of the operator. Thus we can compute all correlators in zero dimensions. In this zero dimensional model, each two point function is just a number. An operator with a normalized two point function is one for which this number is one. This map between two point functions and the norm of states is of course nothing but the usual state operator correspondence.

The dilatation operator allows the Z s to hop between sites of the Cuntz chain. At leading order in large $M + N$, the action of $D_{2\text{eff}}$ is given entirely by $D_{2,0\text{eff}}$. Using the correspondence given in the last equation of Appendix A, the operator equation (2.21) can be translated into an equation for the action of $D_{2\text{eff}}$ on the Cuntz lattice. If all Cuntz lattice occupation numbers are positive then, since $D_{2\text{eff}}$ lowers each occupation number by at most 1, acting with $D_{2\text{eff}}$ can't change the value $p_- = 0$ where p_- is negative the sum of all the negative occupation numbers. This implies that all gauge invariant operators have the same leading two point function and hence, when acting on a Cuntz lattice with, for example, two sites ($\lambda = g^2 N$)

$$g^2 D_{2,0\text{eff}} |\{p_1, p_2\}\rangle = -4\lambda \frac{N+M}{N} (|\{p_1+1, p_2-1\}\rangle - 2|\{p_1, p_2\}\rangle + |\{p_1-1, p_2+1\}\rangle). \quad (2.29)$$

Similarly, if all Cuntz lattice occupation numbers are negative then, since $D_{2\text{eff}}$ raises each occupation number by at most 1, acting with $D_{2\text{eff}}$ again can't change the value $p_- = \sum_i p_i$. Again all gauge invariant operators have the same leading two point function and hence, when acting on a Cuntz lattice with, for example, two sites

$$g^2 D_{2,0\text{eff}} |\{p_1, p_2\}\rangle = -4\lambda \frac{M}{N} (|\{p_1+1, p_2-1\}\rangle - 2|\{p_1, p_2\}\rangle + |\{p_1-1, p_2+1\}\rangle). \quad (2.30)$$

Finally, consider for example the case that $D_{2,0\text{eff}}$ acts on a lattice with two sites and occupation numbers $p_1 = 0, p_2 = 2$. In this case, the normalization of the gauge

invariant operators is not the same: three terms have $p_- = 0$ and one has $p_- = 1$. Taking this into account gives

$$g^2 D_{2,0 \text{ eff}} |\{0, 2\}\rangle = -4\lambda \frac{M+N}{N} (|\{1, 1\}\rangle - |\{0, 2\}\rangle) - 4\lambda \frac{M}{N} \left(\frac{\sqrt{M+N}}{\sqrt{M}} |\{-1, 3\}\rangle - |\{0, 2\}\rangle \right). \quad (2.31)$$

There is a nice convenient way to summarize these results[33, 19, 21]. We will introduce Cuntz oscillators which satisfy the algebra (we associate one of these oscillators to each site of the chain)

$$a^\dagger a = \frac{M}{N} + \theta(\hat{n} + 1) - |0\rangle\langle 0|, \quad aa^\dagger = \frac{M}{N} + \theta(\hat{n} + 1), \quad (2.32)$$

with $\hat{n}$ the number operator. Notice that when $M = 0$ we have only positive occupation numbers so that the above relations reduce to the usual ones

$$a^\dagger a = 1 - |0\rangle\langle 0|, \quad aa^\dagger = 1. \quad (2.33)$$

We can also define these oscillators by giving their action on states of good particle number

$$a|n\rangle = \sqrt{1 + \frac{M}{N}} |n-1\rangle \quad n > 0 \quad (2.34)$$

$$a|n\rangle = \sqrt{\frac{M}{N}} |n-1\rangle \quad n \leq 0. \quad (2.35)$$

In terms of these Cuntz oscillators we have

$$g^2 D_{2 \text{ eff}} = 2\lambda \sum_{l=1}^L (a_l^\dagger - a_{l+1}^\dagger)(a_l - a_{l+1}). \quad (2.36)$$

There is a rather direct way to extract (part of the) geometry of the dual LLM solution from this Cuntz oscillator description[33, 19, 21]. To see this, consider the coherent state

$$|z\rangle = \sum_{n=-\infty}^0 \left(\frac{N}{M}\right)^{\frac{n}{2}} z^n |n\rangle + \sum_{n=1}^{\infty} \left(\frac{N}{M+N}\right)^{\frac{n}{2}} z^n |n\rangle. \quad (2.37)$$

The norm of this state

$$\langle z|z\rangle = \sum_{n=0}^{\infty} \frac{M^n}{N^n |z|^{2n}} + \sum_{n=1}^{\infty} \frac{N^n |z|^{2n}}{(M+N)^n} \quad (2.38)$$

is only finite if $\frac{M}{N} \leq |z|^2 \leq \frac{M+N}{N}$, that is, $|z|^2$ must lie within the annulus. Clearly z is a complex coordinate for the LLM plane.

This one loop result is intriguing: the effect of the background has been completely accounted for by simply modifying the Cuntz oscillator algebra. This is a remarkably simple change. It is natural to ask

Is the net effect of the background (even at higher loops) completely accounted for, by simply modifying the Cuntz oscillator algebra?

We do not have a complete answer to this question. Computing the form of the two loop answer ($D_{4 \text{ eff}}$) is already a rather involved task. We have however studied this question for two sites. In this case

$$g^4 D_{4 \text{ eff}} = -\frac{1}{4}(g^2 D_{2 \text{ eff}})^2 + g^4 \delta D_{4 \text{ eff}} \quad (2.39)$$

where

$$g^4 \delta D_{4 \text{ eff}} = 4\lambda^2 \sum_{l=1}^2 \left(a_l^\dagger [a_{l+1}, a_{l+1}^\dagger] a_{l+1} + a_l^\dagger a_{l+1} [a_l, a_l^\dagger] \right. \\ \left. - a_l^\dagger a_l [a_{l+1}, a_{l+1}^\dagger] - a_l^\dagger [a_l, a_l^\dagger] a_l \right) . \quad (2.40)$$

Thus, at two loops for a Cuntz chain with two sites we find that, again, the net effect of the background is to modify the Cuntz oscillator algebra.

3. Conservation Laws

In the last section we have explained how to extract an effective dilatation operator. Diagonalizing this effective dilatation operator will give the spectrum of anomalous dimensions for a class of operators whose classical dimension is of order N^2 . Non-planar diagrams have to be included to obtain this dilatation operator. One consequence of this is that a spin chain description is no longer useful and we have instead passed to a Cuntz lattice description. In this section we want to answer two questions:

- The Cuntz lattice description can be employed even before a background is introduced. What is the translation of the conserved quantities of the original spin chain into the Cuntz lattice language?
- Is there any evidence that the effective dilatation operator obtained in the presence of the LLM annulus background is integrable?

3.1 U_2 rewritten in the Cuntz Oscillator Language

Before we obtain the conserved charge U_2 in the Cuntz oscillator language, its useful to first write it as a differential operator acting on gauge invariant operators. We will use this expression when we search for a corresponding conserved charge in the nontrivial background. It is a straight forward exercise to find that the planar action of

$$\begin{aligned} U_2 = & \text{Tr} \left((YZZ - ZYZ) \frac{d}{dY} \frac{d}{dZ} \frac{d}{dZ} \right) + \text{Tr} \left((ZZY - YZZ) \frac{d}{dZ} \frac{d}{dY} \frac{d}{dZ} \right) \\ & + \text{Tr} \left((ZYZ - ZZY) \frac{d}{dZ} \frac{d}{dZ} \frac{d}{dY} \right) + \text{Tr} \left((ZYY - YZY) \frac{d}{dZ} \frac{d}{dY} \frac{d}{dY} \right) \\ & + \text{Tr} \left((YYZ - ZYY) \frac{d}{dY} \frac{d}{dZ} \frac{d}{dY} \right) + \text{Tr} \left((YZY - YYZ) \frac{d}{dY} \frac{d}{dY} \frac{d}{dZ} \right), \end{aligned} \quad (3.1)$$

on single trace gauge invariant operators matches the action of U_2 on the spin chain. To illustrate how to obtain a Cuntz oscillator representation, consider the first term above: it acts as

$$ZZY \rightarrow YZZ - ZYZ. \quad (3.2)$$

This term can be represented in terms of Cuntz oscillators as

$$\sum_{l=1}^L (a_{l+1}^\dagger a_{l+1}^\dagger - a_l^\dagger a_{l+1}^\dagger) a_l a_l. \quad (3.3)$$

The second term in the sum should not be truncated to $a_{l+1}^\dagger a_l$ since we have to make sure that there are at least 2 Zs in lattice site l . The result for U_2 is

$$\begin{aligned} U_2 = & \sum_{l=1}^L \left((a_{l+1}^\dagger a_{l+1}^\dagger - a_l^\dagger a_{l+1}^\dagger) a_l a_l + (a_l^\dagger a_l^\dagger - a_{l+1}^\dagger a_{l+1}^\dagger) a_{l+1} a_l \right. \\ & + (a_l^\dagger a_{l-1}^\dagger - a_{l-1}^\dagger a_{l-1}^\dagger) a_l a_l + (a_{l-1}^\dagger - a_l^\dagger) a_{l+1} [a_l, a_l^\dagger] \\ & \left. + (a_{l+1}^\dagger - a_{l-1}^\dagger) [a_l, a_l^\dagger] a_l + (a_l^\dagger - a_{l+1}^\dagger) a_{l-1} [a_l, a_l^\dagger] \right). \end{aligned} \quad (3.4)$$

Using this procedure it is straight forward to write down the Cuntz oscillator representation for any of the conserved charges of the spin chain.

It is tedious but straight forward to compute the commutator of U_2 given above and $D_{2\text{eff}}$ - they do not commute. It seems natural to consider the operator

$$U_{2\text{eff}} = \frac{1}{\chi_B(Z)} U_2 \chi_B(Z). \quad (3.5)$$

It is a simple matter to compute $U_{2\text{eff}}$ using the above expression for U_2 as a differential operator acting on gauge invariant operators. Again, $D_{2\text{eff}}$ and $U_{2\text{eff}}$ do not commute. However, in the leading large M limit the two do commute suggesting that this may be an interesting limit of $D_{2\text{eff}}$. This is explored in detail in section 3.3 below.

3.2 Classical Limit

The original spin chain description of the dilatation operator can be replaced by a sigma model in the limit of a large number of sites. This sigma model precisely matches the Polyakov action describing the propagation of closed strings in $\text{AdS}_5 \times S^5$, in a particular limit[34, 35, 36]. This has also been extended to other examples of the AdS/CFT correspondence[26, 37, 38, 39, 40]. The Cuntz oscillator description of the dilatation operator is simply an alternative language in the undeformed background; when we consider the deformed background the spin chain description is not convenient, so that in this case it is best to use the Cuntz oscillator description. We can obtain a semiclassical description of the Cuntz chain by again considering a large number of sites[40]. In this subsection we will provide further insight into the relation between the spin chain and Cuntz oscillator descriptions by showing that in the dual string theory the two descriptions are simply related by a change of worldsheet gauge choice.

The semi-classical limit of the Cuntz chain is obtained by taking $L \sim \sqrt{N} \rightarrow \infty$, $\lambda \rightarrow \infty$ holding $\frac{\lambda}{L^2}$ fixed and by putting each lattice site into a coherent state (we are discussing the undeformed theory so there are no negatively occupied sites)

$$|z\rangle = \sqrt{1 - |z|^2} \sum_{n=0}^{\infty} z^n |n\rangle, \quad |n\rangle = (a^\dagger)^n |0\rangle. \quad (3.6)$$

The coherent state parameter of the l^{th} site is traded for a radius and an angle $z_l = r_l e^{i\phi_l}$. The action is given, as usual, by

$$S = \int dt \left(i \langle Z | \frac{d}{dt} | Z \rangle - \langle Z | D | Z \rangle \right) \quad |Z\rangle = \prod_l |z_l\rangle. \quad (3.7)$$

After trading the sum over l for an integral over σ , the action is

$$S = -L \int dt \int_0^1 d\sigma \left(\frac{r^2 \dot{\phi}}{1 - r^2} + \frac{\lambda}{L^2} (r'^2 + r^2 \phi'^2) \right). \quad (3.8)$$

For a detailed derivation the reader could consult [40]. We would like to show how this Cuntz chain sigma model can be recovered from the standard string sigma model action on $R \times S^3$.

A string moving on $R \times S^3$ can be described by the principal chiral model with $\text{su}(2)$ valued currents

$$j_\tau = g^{-1} \frac{\partial g}{\partial \tau}, \quad j_\sigma = g^{-1} \frac{\partial g}{\partial \sigma}, \quad (3.9)$$

where

$$g = \begin{bmatrix} Z & iY \\ i\bar{Y} & \bar{Z} \end{bmatrix} \in \text{SU}(2), \quad (3.10)$$

and Z and Y are the coordinates of a sphere

$$|Z|^2 + |Y|^2 = 1. \quad (3.11)$$

We are choosing to employ a principal chiral model description since this description manifests the integrability of the model. Parametrize the sphere coordinates as follows

$$Z = r e^{i(\kappa\tau - \phi)}, \quad Y = \sqrt{1 - r^2} e^{i(\varphi + \kappa\tau)}. \quad (3.12)$$

The equations of motion

$$\partial_\tau j_\tau - \partial_\sigma j_\sigma = 0 \quad (3.13)$$

can be obtained from the Polyakov action in conformal gauge and after fixing the residual conformal diffeomorphism freedom by choosing $t = \kappa\tau$. In this gauge, energy is homogeneously distributed along σ . To obtain the low energy limit, we take $\kappa \rightarrow \infty$ holding $\kappa\dot{r}$, $\kappa\dot{\phi}$ and $\kappa\dot{\varphi}$ fixed. It is precisely in this gauge and in this limit that [34] matched the semiclassical limit of the one loop spin chain to the string sigma model. The Lagrangian in this limit becomes

$$\mathcal{L} = -\frac{1}{4}(j_\tau^2 - j_\sigma^2) = r^2 \kappa \dot{\phi} - (1 - r^2) \kappa \dot{\varphi} + \frac{1}{2} r^2 \phi'^2 + \frac{1}{2} (1 - r^2) \varphi'^2 + \frac{1}{2} \frac{r'^2}{1 - r^2}.$$

The equations of motion following from this action needs to be supplemented by the usual Virasoro constraints, which in this limit are $j_\tau^2 + j_\sigma^2 = 0$ and

$$\kappa \varphi' (1 - r^2)^2 + \kappa \phi' r^2 (r^2 - 1) + O(1) = 0. \quad (3.14)$$

The Cuntz sigma model (3.8) does not contain the field φ . Thus, it should be eliminated before we can expect to obtain agreement. Integrate by parts to obtain

$$\mathcal{L} = r^2 \kappa \dot{\phi} - \varphi \frac{\partial}{\partial \tau} (1 - r^2) \kappa + \frac{1}{2} r^2 \phi'^2 + \frac{1}{2} (1 - r^2) \varphi'^2 + \frac{1}{2} \frac{r'^2}{1 - r^2}. \quad (3.15)$$

Now, using the φ equation of motion (to rewrite the coefficients of φ in the action) and the (square of the) Virasoro constraint (3.14) we find

$$\mathcal{L} = r^2 \kappa \dot{\phi} + \frac{1}{2} r^2 \phi'^2 + \frac{1}{2} \frac{r'^2}{1 - r^2} - \frac{1}{2} \frac{r^4}{1 - r^2} \phi'^2. \quad (3.16)$$

This does not agree with the result (3.8).

The disagreement is not surprising: the sigma model (3.16) is written in a gauge in which energy is distributed homogeneously along the string; the sigma model (3.8) corresponds to a gauge in which p_φ (= angular momentum conjugate to φ) is distributed

homogeneously along the string. In the Cuntz chain, only Y s mark lattice sites; in the usual spin chain both Y s and Z s mark lattice sites. Consequently to go from the σ_{cc} coordinate of the Cuntz chain to the σ_{sc} coordinate of the spin chain, we need to “add the Z s back in”

$$\sigma_{sc} = \sigma_{cc} + \int_0^{\sigma_{cc}} n_z(\sigma') d\sigma' = \sigma_{cc} + \int_0^{\sigma_{cc}} \frac{r^2}{1-r^2} d\sigma', \quad \tau_{sc} = \tau_{cc}. \quad (3.17)$$

In this last equation, $n_z(\sigma') = r^2/(1-r^2)$ is the expected number of Cuntz particles (= number of Z s) at σ' . It is now straight forward to compute

$$\frac{\partial \sigma_{sc}}{\partial \sigma_{cc}} = \frac{1}{1-r^2}, \quad \frac{\partial \tau_{sc}}{\partial \tau_{cc}} = 1, \quad \frac{\partial \tau_{sc}}{\partial \sigma_{cc}} = 0, \quad (3.18)$$

$$\frac{\partial \sigma_{sc}}{\partial \tau_{cc}} = -2r^2 \frac{\partial \phi}{\partial \sigma_{cc}} = -2 \frac{r^2}{1-r^2} \frac{\partial \phi}{\partial \sigma_{sc}}. \quad (3.19)$$

The integrability condition

$$\frac{\partial}{\partial \tau_{cc}} \frac{\partial}{\partial \sigma_{cc}} \sigma_{sc} = \frac{\partial}{\partial \tau_{cc}} \frac{1}{1-r^2} = \frac{\partial}{\partial \sigma_{cc}} \frac{\partial}{\partial \tau_{cc}} \sigma_{sc} = \frac{\partial}{\partial \sigma_{cc}} \left(-2r^2 \frac{\partial \phi}{\partial \sigma_{cc}} \right) \quad (3.20)$$

is nothing but the ϕ equation of motion derived from (3.8). It is now a straight forward exercise to verify that after this change of coordinates (3.8) and (3.16) match perfectly.

The Cuntz oscillator Hamiltonian (3.8) can be written as

$$S = -L \int dt \int_0^1 d\sigma \left(n_z(\sigma) \dot{\phi} + \frac{\lambda}{L^2} (r'^2 + r^2 \phi'^2) \right). \quad (3.21)$$

The advantage of this rewriting is that it holds in general - that is, both for the undeformed and deformed backgrounds. Inserting the explicit expression for the expected number of Cuntz particles in the deformed background we find

$$S = -L \int dt \int_0^1 d\sigma \left(\left[\frac{r^2}{1 + \frac{M}{N} - r^2} - \frac{\frac{M}{N}}{\frac{M}{N} - r^2} \right] \dot{\phi} + \frac{\lambda}{L^2} (r'^2 + r^2 \phi'^2) \right). \quad (3.22)$$

If one drops either of the two terms in square braces one obtains a model which can be related to the low energy limit of the principal chiral model and hence is the low energy limit of an integrable model. The physical interpretation of such a truncation is clear: keeping only the first term corresponds to focusing on fluctuations localized at the outer edge of the annulus; keeping only the second term corresponds to focusing on fluctuations localized at the inner edge of the annulus. For these classes of fluctuations, it seems that the dynamics is integrable. It would be interesting to establish if the full model is integrable or not.

There is a very natural generalization to multi ring LLM geometries, corresponding to backgrounds created by Schur polynomials labeled by a Young diagram with more than 4 edges and each edge with a length of $O(N)$. In this case $n_z(\sigma)$ is a sum of terms, one for each edge. Restricting to fluctuations localized about a particular edge again gives a model which can be related to the low energy limit of the principal chiral model and hence is the low energy limit of an integrable model. These localized excitations have been constructed in [19].

The superstar geometry[20] has been related to an LLM geometry with boundary condition given by a sequence of concentric alternating black and white rings[30]. Rings of the same color have the same area and the total area of the black rings is π . As mentioned above, to construct $n_z(\sigma)$ we need to sum a term for each edge of the multi-ring geometry. We will consider a geometry which corresponds to the Young diagram shown in figure 1 with $n_1, n_2 \ll N$. In this case, we sum a very large number of terms and hence may use the Euler-Maclaurin formula to rewrite the sum as an integral. Carrying this integral out we find (we dropped an additive constant that will not contribute to the equations of motion)

$$n_z(\sigma) = \frac{\alpha}{\alpha + \beta} \frac{r^2}{1 + \frac{M}{N} - r^2} \quad (3.23)$$

where

$$\alpha = \frac{n_1}{N}, \quad \beta = \frac{n_2}{N}. \quad (3.24)$$




Figure 1: The Young diagram corresponding to the superstar geometry.

Thus, the semiclassical limit of the Cuntz chain model is

$$S = -L \int dt \int_0^1 d\sigma \left(\frac{\alpha}{\alpha + \beta} \frac{r^2}{1 + \frac{M}{N} - r^2} \dot{\phi} + \frac{\lambda}{L^2} (r'^2 + r^2 \phi'^2) \right). \quad (3.25)$$

After rescaling t we can recover the action (3.8) up to an overall multiplicative constant. Thus, the model can again be related to the low energy limit of an integrable model.

3.3 Integrability in the Large M Limit

In this subsection we will consider the large M limit, that is, we take $M, N \rightarrow \infty$ and in addition, the ratio $\frac{N}{M} \rightarrow 0$. In this limit we suppress all $\frac{N}{M}$ dependence. The dilatation operator for the sector we consider, after subtracting the classical dimension out, can be written as

$$D = D \left(Z, Y, \frac{d}{dZ}, \frac{d}{dY} \right). \quad (3.26)$$

To get the large M limit of D_{eff} (we denote this operator by $\tilde{D}_{\text{eff}}$) we should simply replace $\frac{d}{dZ}$ by MZ^{-1} in the above expression to obtain

$$\tilde{D}_{\text{eff}} = \tilde{D}_{\text{eff}} \left(Z, Y, MZ^{-1}, \frac{d}{dY} \right). \quad (3.27)$$

Expand this operator as

$$\tilde{D}_{\text{eff}} = \sum_n \tilde{D}_{\text{eff } n} \quad (3.28)$$

where $\tilde{D}_{\text{eff } n}$ has a total of n derivatives with respect to Y . From the general structure of a connected planar l -loop vertex we know that D will act on $l+1$ adjacent sites; thus, it will contain $l+1$ derivatives. The leading contribution in the large M limit would come from terms which have all $l+1$ derivatives acting on Z and these are replaced to give an $M^{l+1} \text{Tr}(Z^{-l-1})$. This leading term is captured in $\tilde{D}_{\text{eff } 0}$, which up to an arbitrary coefficient, is now determined. Since the dimension of $\text{Tr}(Z^J)$ is not corrected, it must be annihilated by D and hence, in the large M limit it must be annihilated by $\tilde{D}_{\text{eff } 0}$. This implies that $\tilde{D}_{\text{eff } 0}$ vanishes. Thus, the leading contribution will in fact come from the $\tilde{D}_{\text{eff } 1}$. The l -loop term will thus have a $(g^2 M)^l$ dependence - this is the dependence that the present argument captures². Since D is dimensionless, and preserves the total number of Z s and the total number of Y s, it is clear that, to leading order at large M

$$\tilde{D}_{\text{eff}} = \tilde{D}_{\text{eff } 1} = \sum_n c_n \text{Tr} \left(Z^n Y Z^{-n} \frac{d}{dY} \right) \quad (3.29)$$

where the c_n depend on $g^2 M$. It is trivial to see that

$$\left[\text{Tr} \left(Z^{-n} Y Z^n \frac{d}{dY} \right), \text{Tr} \left(Z^{-m} Y Z^m \frac{d}{dY} \right) \right] = 0, \quad (3.30)$$

²Of course, our large M limit is a double scaling limit in which we take $M \rightarrow \infty$, $g^2 \rightarrow 0$ holding $g^2 M$ fixed. This limit is the natural one: our effective genus counting parameter is $\frac{1}{M}$ so that $\lambda = g^2 M$ is the obvious definition of the 't Hooft coupling. See [22] for further details.

so that

$$\left[\tilde{D}_{\text{eff}}, \text{Tr} \left(Z^{-m} Y Z^m \frac{d}{dY} \right) \right] = 0, \quad (3.31)$$

which clearly demonstrates an infinite number of conserved quantities to all loops.

What is the physical meaning of this limit? Recall that the dilatation operator can be read from the two point functions of the theory. Restricting to operators constructed from scalar fields only is, in general, not possible due to operator mixing. However, it is possible to show[3] that it is consistent to restrict to operators built using only the two complex fields Y and Z . In this case, we can compute the two point functions in a reduced model comprising of only the Y and Z matrices. Interpreted in this way, the dilatation operator can be understood as implementing the Wick contractions associated with the F-term vertex. For example, consider the combination

$$\frac{d}{dZ} + MZ^{-1} \quad (3.32)$$

which replaces $\frac{d}{dZ}$ in transforming the undeformed into the deformed Cuntz chain. The $\frac{d}{dZ}$ term represents a contraction between the vertex and a $Z^\dagger$ in one of the fields whose two point function we are computing; to see this connection it is useful to remember that

$$\langle Z_{ij}^\dagger Z_{kl} \rangle = \delta_{jk} \delta_{il} = \frac{d}{dZ_{ji}} Z_{kl}. \quad (3.33)$$

In contrast to this, the MZ^{-1} term represents a contraction between the vertex and a $Z^\dagger$ in the operator representing the background. In the large M limit, the contractions with the background completely dominate as compared to contractions with fields belonging to the operators of the two point functions we are computing. One can think that the matrices entering into the operators are “bits of a string”. In the limit that we consider, the different bits in the string do not interact with each other - they interact only with the background. We would indeed expect the dynamics to simplify in this limit.

In the large M limit, the action of the Cuntz chain (3.22) becomes

$$S = -L \int dt \int_0^1 d\sigma \left(-\dot{\phi} + \frac{\lambda}{L^2} (r'^2 + r^2 \phi'^2) \right). \quad (3.34)$$

Since $\dot{\phi}$ is a total derivative, all time derivatives drop out of the equations of motion. This implies that the dynamics becomes trivial which is indeed consistent with integrability. It is rather interesting that there is a class of operators in $\mathcal{N} = 4$ super Yang-Mills theory that have such a simple description.

In terms of the dual LLM boundary condition, the large M limit corresponds to an annulus with a large radius and fixed area, so that the annulus is becoming very thin.

4. Discussion

The problem of computing the anomalous dimensions of operators with a large ($O(N^2)$) R-charge corresponds to a generalization, in the dual gravitational description, to string dynamics in spacetimes that are only asymptotically $\text{AdS}_5 \times \text{S}^5$. The problem can again be reduced to diagonalizing a Hamiltonian. In the $\text{AdS}_5 \times \text{S}^5$ spacetime, this Hamiltonian was an integrable spin chain. As a consequence of the fact that our strings can exchange angular momentum with the background, our Hamiltonian describes Cuntz particles hopping on a lattice. In the gauge theory description, the terms in the dilatation operator that allow the strings to exchange angular momentum with the background arise from summing (an infinite number of) non-planar corrections. It is surprisingly straight forward to write down very explicit expressions for the relevant Cuntz chain Hamiltonians.

A natural question to ask is if our Cuntz chain Hamiltonians correspond to integrable systems. We don't know. However, we have given some evidence that the large M limit of our Hamiltonian does admit higher conserved charges and that certain localized semiclassical excitations are described by the low energy limit of a principal chiral model, so an optimist would indeed conjecture that our Cuntz chain Hamiltonian is integrable. We hope that we have managed to convince the reader (even if she is pessimistic) that these are interesting limits of the original $\mathcal{N} = 4$ super Yang-Mills theory that warrant further study.

Acknowledgements: We would like to thank Tom Brown, Kevin Goldstein, Vishnu Jejjala, Yusuke Kimura, Jeff Murugan and Sanjaye Ramgoolam for enjoyable, helpful discussions. This work is based upon research supported by the South African Research Chairs Initiative of the Department of Science and Technology and National Research Foundation. Any opinion, findings and conclusions or recommendations expressed in this material are those of the authors and therefore the NRF and DST do not accept any liability with regard thereto. This work is also supported by NRF grant number Gun 2047219.

A. Two Point Functions

In this Appendix we will compute the two point functions used in section 2. We only want these two point functions to the leading order in an $M + N$ expansion. We will use f_B to denote the product of the weights of Young diagram B . It is straight forward to obtain

$$f_B = \frac{G_2(N + M + 1)}{G_2(N + 1)G_2(M + 1)}, \quad (\text{A.1})$$

where $G_2(n+1)$ is the Barnes function defined by ($\Gamma(z)$ is the Gamma function)

$$G_2(z+1) = \Gamma(z)G_2(z). \quad (\text{A.2})$$

In particular, for an integer $z = n$ we have

$$G_2(n+1) = \prod_{k=1}^{n-1} k!. \quad (\text{A.3})$$

First consider the free field theory two point function

$$\langle \chi_B(Z) \chi_B(Z^\dagger) \text{Tr}(Y^2 Z^{J_0}) \text{Tr}(Y^2 Z^{J_0})^\dagger \rangle. \quad (\text{A.4})$$

According to [21], this two point function is given, at the leading order in a large $M+N$ expansion, by f_B times $\left(\frac{M+N}{N}\right)^{J_0}$ times the free field theory two point function in the trivial background

$$\langle \text{Tr}(Y^2 Z^{J_0}) \text{Tr}(Y^2 Z^{J_0})^\dagger \rangle = N^{J_0+2} + O(N^{J_0}). \quad (\text{A.5})$$

Thus,

$$\langle \chi_B(Z) \chi_B(Z^\dagger) \text{Tr}(Y^2 Z^{J_0}) \text{Tr}(Y^2 Z^{J_0})^\dagger \rangle = f_B (N+M)^{J_0} N^2. \quad (\text{A.6})$$

Arguing in exactly the same way, we find

$$\langle \chi_B(Z) \chi_B(Z^\dagger) \text{Tr}(Y Z^n Y Z^{J_0-n}) \text{Tr}(Y Z^m Y Z^{J_0-m})^\dagger \rangle = f_B (N+M)^{J_0} N^2 \delta_{mn}. \quad (\text{A.7})$$

Next, consider

$$\left\langle \chi_{B,B'}^{(1)}(Z, Y Z^{J_0+1} Y) \chi_{B,B'}^{(1)}(Z, Y Z^{J_0+1} Y)^\dagger \right\rangle \quad (\text{A.8})$$

where

$$\chi_{B,B'}^{(1)}(Z, Y Z^{J_0+1} Y) = (Y Z^{J_0+1} Y)_{ij} \frac{\partial}{\partial Z_{ij}} \chi_B(Z). \quad (\text{A.9})$$

These correlators have been computed in [41]. The result is

$$\left\langle \chi_{B,B'}^{(1)}(Z, W) \chi_{B,B'}^{(1)}(Z^\dagger, W^\dagger) \right\rangle = \frac{\text{hooks}_B}{\text{hooks}_{B'}} f_B F_0 + c_{BB'} f_B F_1. \quad (\text{A.10})$$

We find $c_{BB'} = M$ and $\frac{\text{hooks}_B}{\text{hooks}_{B'}} = MN$. Further, for the open string word $W = Y Z^{J_0+1} Y$ we find that at the leading order

$$F_0 = N^{J_0+2} \left(\frac{M+N}{N} \right)^{J_0+1}, \quad F_1 \sim N^{J_0-1} \left(\frac{M+N}{N} \right)^{J_0+1}, \quad (\text{A.11})$$

so that, to the leading order we have

$$\left\langle \chi_{B,B'}^{(1)}(Z, YZ^{J_0+1}Y) \chi_{B,B'}^{(1)}(Z, YZ^{J_0+1}Y)^\dagger \right\rangle = f_B M N^2 (M+N)^{J_0+1}. \quad (\text{A.12})$$

This correlator result can also be written as

$$\left\langle \chi_B(Z) \chi_B(Z^\dagger) \text{Tr}(YZ^{-1}YZ^{J_0+1}) \text{Tr}(YZ^{-1}YZ^{J_0+1})^\dagger \right\rangle = f_B \frac{1}{M} (N+M)^{J_0+1} N^2. \quad (\text{A.13})$$

Again arguing in exactly the same way, we find

$$\left\langle \chi_B(Z) \chi_B(Z^\dagger) \text{Tr}(YZ^{-n}YZ^{J_0+n}) \text{Tr}(YZ^{-m}YZ^{J_0+m})^\dagger \right\rangle = f_B \frac{1}{M^n} (N+M)^{J_0+n} N^2 \delta_{mn}. \quad (\text{A.14})$$

Notice that at large enough M that $\frac{N}{M}$ can be neglected, all the gauge invariant operators considered have exactly the same two point function.

Finally, by using the methods of this Appendix, we can obtain the general result

$$\langle \mathcal{O}_B(\{p\}, J_0) \mathcal{O}_B(\{p\}, J_0)^\dagger \rangle = N^2 f_B \frac{(M+N)^{J_0+p_-}}{M^{p_-}}, \quad (\text{A.15})$$

where $\{p\}$ denotes the occupation numbers of the Cuntz chain and p_- is negative the sum of all the negative occupation numbers. Thus, we have the correspondence

$$\mathcal{O}_B(\{p\}, J_0) \leftrightarrow \sqrt{N^2 f_B \frac{(M+N)^{J_0+p_-}}{M^{p_-}}} |\{p\}\rangle, \quad (\text{A.16})$$

between operators and normalized Cuntz lattice states.

B. More on the Cuntz Chain

To specify the general Cuntz chain model (3.21) one needs to specify the expected number of Cuntz particles $n_z(\sigma)$. Given $n_z(\sigma)$, what is the corresponding supergravity background? Using the results of the first of [33] as well as (3.21), the metric on the $y=0$ plane and the circle along which the string moves (parametrized by³ φ) can be written as

$$ds^2 = -h^{-2}(Dt)^2 + h^2 dz d\bar{z} + h^{-2} d\varphi^2, \quad Dt = dt - \frac{1}{2} i \bar{V} dz + \frac{1}{2} i V d\bar{z}, \quad (\text{B.1})$$

$$V = \frac{n_z}{\bar{z}}, \quad h^4 = \frac{\partial V}{\partial z}, \quad z = r e^{i\phi}. \quad (\text{B.2})$$

³The angular momentum along this circle is due to the Y fields appearing in the gauge invariant operator dual to the string.

C. Explicit Expressions for the Two Loop Dilatation Operator

In these expressions hatted indices are again to be dropped, $\theta(x) = 1$ if $x > 0$ and vanishes otherwise. To obtain this result we have assumed $J_0 > 0$, $J_0 - p \geq 0$ and $p \geq 0$ - assumptions which can easily be relaxed if need be.

$$\begin{aligned} \delta D_{4,0 \text{ eff}} \mathcal{O}_p^{J_0; J_1, \dots, J_k} &= 4(\mathcal{O}_1^{J_0; J_1, \dots, J_k} - \mathcal{O}_0^{J_0; J_1, \dots, J_k}) \times \\ &\times N(N+M)(\delta_{p=0} + \delta_{p=J_0} - \delta_{p=1} - \delta_{p=J_0-1}), \end{aligned} \quad (\text{C.1})$$

$$\begin{aligned} \delta D_{4,+ \text{ eff}} \mathcal{O}_p^{J_0; J_1, \dots, J_k} &= 4(M+2N)\theta(p-1) \times \\ &\times (\mathcal{O}_0^{J_0-p+1; J_1, \dots, J_k, p-1} - \mathcal{O}_1^{J_0-p+1; J_1, \dots, J_k, p-1}) \\ &+ 4(M+2N)\theta(J_0-p-1)(\mathcal{O}_0^{p+1; J_1, \dots, J_k, J_0-p-1} - \mathcal{O}_1^{p+1; J_1, \dots, J_k, J_0-p-1}) \\ &+ 4(N+M)\theta(p)(\mathcal{O}_1^{J_0-p; J_1, \dots, J_k, p} - \mathcal{O}_0^{J_0-p; J_1, \dots, J_k, p}) \\ &+ 4(N+M)\theta(J_0-p)(\mathcal{O}_1^{p; J_1, \dots, J_k, J_0-p} - \mathcal{O}_0^{p; J_1, \dots, J_k, J_0-p}) \\ &+ \sum_{s=1}^{J_0-1} 4N(\delta_{p=0} + \delta_{p=J_0})(\mathcal{O}_1^{J_0-s; J_1, \dots, J_k, s} - \mathcal{O}_0^{J_0-s; J_1, \dots, J_k, s}), \end{aligned} \quad (\text{C.2})$$

$$\begin{aligned} \delta D_{4,- \text{ eff}} \mathcal{O}_p^{J_0; J_1, \dots, J_k} &= 4N \sum_{j=1}^k J_j (\delta_{p=0} + \delta_{p=J_0}) \times \\ &\times (\mathcal{O}_1^{J_0+J_j; J_1, \dots, \hat{J}_j, \dots, J_k} - \mathcal{O}_0^{J_0+J_j; J_1, \dots, \hat{J}_j, \dots, J_k}) \\ &- 4N(\delta_{p=0} + \delta_{p=J_0}) \sum_{i=1}^k \delta_{J_i=1} (\mathcal{O}_1^{J_0+1; J_1, \dots, \hat{J}_i, \dots, J_k} - \mathcal{O}_0^{J_0+1; J_1, \dots, \hat{J}_i, \dots, J_k}), \end{aligned} \quad (\text{C.3})$$

$$\begin{aligned} \delta D_{4,+ - \text{ eff}} \mathcal{O}_p^{J_0; J_1, \dots, J_k} &= 4\theta(p) \sum_{i=1}^k J_i (\mathcal{O}_1^{J_0+J_i-p; J_1, \dots, \hat{J}_i, \dots, J_k, p} \\ &- \mathcal{O}_0^{J_0+J_i-p; J_1, \dots, \hat{J}_i, \dots, J_k, p}) + 4\theta(J_0-p) \sum_{i=1}^k J_i (\mathcal{O}_1^{J_i+p; J_1, \dots, \hat{J}_i, \dots, J_k, J_0-p} \\ &- \mathcal{O}_0^{J_i+p; J_1, \dots, \hat{J}_i, \dots, J_k, J_0-p}) - 4\theta(J_i-1) \sum_{i=1}^k J_i (\mathcal{O}_1^{p+1; J_1, \dots, \hat{J}_i, \dots, J_k, J_0+J_i-p-1} \\ &- \mathcal{O}_0^{p+1; J_1, \dots, \hat{J}_i, \dots, J_k, J_0+J_i-p-1}) - 4\theta(J_i-1) \sum_{i=1}^k J_i (\mathcal{O}_1^{J_0-p+1; J_1, \dots, \hat{J}_i, \dots, J_k, J_i+p-1} \end{aligned}$$

$$-\mathcal{O}_0^{J_0-p+1;J_1,\dots,\hat{J}_i,\dots,J_k,J_i+p-1}), \quad (\text{C.4})$$

$$\begin{aligned} \delta D_{4,++} \text{ eff } \mathcal{O}_p^{J_0;J_1,\dots,J_k} &= 4 \sum_{r=1}^{p-2} (\mathcal{O}_0^{J_0-p+1;J_1,\dots,J_k,r,p-r-1} \\ &- \mathcal{O}_1^{J_0-p+1;J_1,\dots,J_k,r,p-r-1}) + 4 \sum_{r=1}^{J_0-p-2} (\mathcal{O}_0^{p+1;J_1,\dots,J_k,r,J_0-p-r-1} \\ &- \mathcal{O}_1^{p+1;J_1,\dots,J_k,r,J_0-p-r-1}) + 4\theta(p) \sum_{s=1}^{J_0-p-1} (\mathcal{O}_1^{J_0-p-s;J_1,\dots,J_k,s,p} \\ &- \mathcal{O}_0^{J_0-p-s;J_1,\dots,J_k,s,p}) + 4\theta(J_0-p) \sum_{s=1}^{p-1} (\mathcal{O}_1^{p-s;J_1,\dots,J_k,s,J_0-p} \\ &- \mathcal{O}_0^{p-s;J_1,\dots,J_k,s,J_0-p}), \end{aligned} \quad (\text{C.5})$$

$$\delta D_{4,--} \text{ eff } \mathcal{O}_p^{J_0;J_1,\dots,J_k} = 0. \quad (\text{C.6})$$

Setting $M = 0$ in the above expressions gives exact agreement with Appendix E of [3] except for the last term in our expression for $\delta D_{4,-}$. The extra term that we have ensures that no joinings between the trace with the Y s and a trace without Y s and a single Z can occur. In this case, the $\frac{d}{dZ_{ij}}$ in $\delta D_{4 \text{ eff}}$ acts on $\text{Tr}(Z)$ to produce δ_{ij} . This vanishes because $\frac{d}{dZ_{ij}}$ appears inside a commutator; the extra term is needed.

References

- [1] J. A. Minahan and K. Zarembo, “The Bethe-ansatz for $N = 4$ super Yang-Mills,” JHEP **0303**, 013 (2003) [arXiv:hep-th/0212208].
- [2] N. Beisert and M. Staudacher, “The $N=4$ SYM Integrable Super Spin Chain,” Nucl. Phys. B **670**, 439 (2003) [arXiv:hep-th/0307042].
- [3] N. Beisert, C. Kristjansen and M. Staudacher, “The dilatation operator of $N = 4$ super Yang-Mills theory,” Nucl. Phys. B **664**, 131 (2003) [arXiv:hep-th/0303060].
- [4] N. Beisert, “The $su(2-3)$ dynamic spin chain,” Nucl. Phys. B **682**, 487 (2004) [arXiv:hep-th/0310252].
- [5] D. Serban and M. Staudacher, “Planar $N = 4$ gauge theory and the Inozemtsev long range spin chain,” JHEP **0406**, 001 (2004) [arXiv:hep-th/0401057].
- [6] N. Beisert, V. Dippel and M. Staudacher, “A novel long range spin chain and planar $N = 4$ super Yang-Mills,” JHEP **0407**, 075 (2004) [arXiv:hep-th/0405001].

- [7] B. Eden, C. Jarczak and E. Sokatchev, “A three-loop test of the dilatation operator in $N = 4$ SYM,” Nucl. Phys. B **712**, 157 (2005) [arXiv:hep-th/0409009].
- [8] M. Staudacher, “The factorized S-matrix of CFT/AdS,” JHEP **0505**, 054 (2005) [arXiv:hep-th/0412188].
- [9] V. A. Kazakov and K. Zarembo, “Classical / quantum integrability in non-compact sector of AdS/CFT,” JHEP **0410**, 060 (2004) [arXiv:hep-th/0410105].
- [10] N. Beisert, “The $su(2-2)$ dynamic S-matrix,” Adv. Theor. Math. Phys. **12**, 945 (2008) [arXiv:hep-th/0511082].
- [11] B. I. Zwiebel, “ $N = 4$ SYM to two loops: Compact expressions for the non-compact symmetry algebra of the $su(1,1-2)$ sector,” JHEP **0602**, 055 (2006) [arXiv:hep-th/0511109].
- [12] N. Beisert and M. Staudacher, “Long-range $PSU(2,2-4)$ Bethe ansatz for gauge theory and strings,” Nucl. Phys. B **727**, 1 (2005) [arXiv:hep-th/0504190].
- [13] N. Beisert, “An $SU(1-1)$ -invariant S-matrix with dynamic representations,” Bulg. J. Phys. **33S1**, 371 (2006) [arXiv:hep-th/0511013].
- [14] B. Eden and M. Staudacher, “Integrability and transcendentality,” J. Stat. Mech. **0611**, P014 (2006) [arXiv:hep-th/0603157].
- [15] N. Beisert, R. Hernandez and E. Lopez, “A crossing-symmetric phase for $AdS(5) \times S^5$ strings,” JHEP **0611**, 070 (2006) [arXiv:hep-th/0609044].
- [16] N. Beisert, B. Eden and M. Staudacher, “Transcendentality and crossing,” J. Stat. Mech. **0701**, P021 (2007) [arXiv:hep-th/0610251].
- [17] C. Kristjansen, M. Orselli and K. Zoubos, “Non-planar ABJM Theory and Integrability,” JHEP **0903**, 037 (2009) [arXiv:0811.2150 [hep-th]],
P. Caputa, C. Kristjansen and K. Zoubos, “Non-planar ABJ Theory and Parity,” Phys. Lett. B **677**, 197 (2009) [arXiv:0903.3354 [hep-th]].
- [18] V. Balasubramanian, M. Berkooz, A. Naqvi and M. J. Strassler, “Giant gravitons in conformal field theory,” JHEP **0204**, 034 (2002) [arXiv:hep-th/0107119].
- [19] R. de Mello Koch, “Geometries from Young Diagrams,” JHEP **0811**, 061 (2008) [arXiv:0806.0685 [hep-th]].
- [20] R. C. Myers and O. Tafjord, “Superstars and giant gravitons,” JHEP **0111**, 009 (2001) [arXiv:hep-th/0109127].
- [21] R. de Mello Koch, N. Ives and M. Stephanou, “Correlators in Nontrivial Backgrounds,” Phys. Rev. D **79**, 026004 (2009) [arXiv:0810.4041 [hep-th]].
- [22] R. de Mello Koch, T. K. Dey, N. Ives and M. Stephanou, “Correlators Of Operators with a Large R-charge,” arXiv:0905.2273 [hep-th].
- [23] S. Corley, A. Jevicki and S. Ramgoolam, “Exact correlators of giant gravitons from dual $N = 4$ SYM theory,” Adv. Theor. Math. Phys. **5**, 809 (2002) [arXiv:hep-th/0111222].

- [24] S. Corley and S. Ramgoolam, “Finite factorization equations and sum rules for BPS correlators in $N = 4$ SYM theory,” Nucl. Phys. B **641**, 131 (2002) [arXiv:hep-th/0205221].
- [25] S. Ramgoolam, “Wilson loops in 2-D Yang-Mills: Euler characters and loop equations,” Int. J. Mod. Phys. A **11**, 3885 (1996) [arXiv:hep-th/9412110],
R. de Mello Koch and R. Gwyn, “Giant graviton correlators from dual $SU(N)$ super Yang-Mills theory,” JHEP **0411**, 081 (2004) [arXiv:hep-th/0410236].
- [26] D. Berenstein, D. H. Correa and S. E. Vazquez, “Quantizing open spin chains with variable length: An example from giant gravitons,” Phys. Rev. Lett. **95**, 191601 (2005) [arXiv:hep-th/0502172].
- [27] Y. Kimura and S. Ramgoolam, “Branes, Anti-Branes and Brauer Algebras in Gauge-Gravity duality,” JHEP **0711**, 078 (2007) [arXiv:0709.2158 [hep-th]],
T. W. Brown, P. J. Heslop and S. Ramgoolam, “Diagonal multi-matrix correlators and BPS operators in $N=4$ SYM,” JHEP **0802**, 030 (2008) [arXiv:0711.0176 [hep-th]],
R. Bhattacharyya, S. Collins and R. d. M. Koch, “Exact Multi-Matrix Correlators,” JHEP **0803**, 044 (2008) [arXiv:0801.2061 [hep-th]],
S. Ramgoolam, “Schur-Weyl duality as an instrument of Gauge-String duality,” AIP Conf. Proc. **1031**, 255 (2008) [arXiv:0804.2764 [hep-th]],
R. Bhattacharyya, R. de Mello Koch and M. Stephanou, “Exact Multi-Restricted Schur Polynomial Correlators,” JHEP **0806**, 101 (2008) [arXiv:0805.3025 [hep-th]],
T. W. Brown, P. J. Heslop and S. Ramgoolam, “Diagonal free field matrix correlators, global symmetries and giant gravitons,” JHEP **0904**, 089 (2009) [arXiv:0806.1911 [hep-th]],
Y. Kimura and S. Ramgoolam, “Enhanced symmetries of gauge theory and resolving the spectrum of local operators,” Phys. Rev. D **78**, 126003 (2008) [arXiv:0807.3696 [hep-th]]. Y. Kimura, “Non-holomorphic multi-matrix gauge invariant operators based on Brauer algebra,” arXiv:0910.2170 [hep-th].
- [28] D. Berenstein, “A toy model for the AdS/CFT correspondence,” JHEP **0407**, 018 (2004) [arXiv:hep-th/0403110].
- [29] H. Lin, O. Lunin and J. M. Maldacena, “Bubbling AdS space and $1/2$ BPS geometries,” JHEP **0410**, 025 (2004) [arXiv:hep-th/0409174].
- [30] V. Balasubramanian, J. de Boer, V. Jejjala and J. Simon, “The library of Babel: On the origin of gravitational thermodynamics,” JHEP **0512**, 006 (2005) [arXiv:hep-th/0508023],
V. Balasubramanian, V. Jejjala and J. Simon, “The library of Babel,” Int. J. Mod. Phys. D **14**, 2181 (2005) [arXiv:hep-th/0505123].
- [31] K. Skenderis and M. Taylor, “Anatomy of bubbling solutions,” JHEP **0709**, 019 (2007) [arXiv:0706.0216 [hep-th]].
- [32] J. Simon, “Small Black holes vs horizonless solutions in AdS,” arXiv:0910.3225 [hep-th].

- [33] S. E. Vazquez, “Reconstructing 1/2 BPS space-time metrics from matrix models and spin chains,” *Phys. Rev. D* **75**, 125012 (2007) [arXiv:hep-th/0612014],
H. Y. Chen, D. H. Correa and G. A. Silva, “Geometry and topology of bubble solutions from gauge theory,” *Phys. Rev. D* **76**, 026003 (2007) [arXiv:hep-th/0703068].
- [34] M. Kruczenski, “Spin chains and string theory,” *Phys. Rev. Lett.* **93**, 161602 (2004) [arXiv:hep-th/0311203],
M. Kruczenski, A. V. Ryzhov and A. A. Tseytlin, “Large spin limit of $\text{AdS}(5) \times S^{*5}$ string theory and low energy expansion of ferromagnetic spin chains,” *Nucl. Phys. B* **692**, 3 (2004) [arXiv:hep-th/0403120].
- [35] R. Hernandez and E. Lopez, “The $\text{SU}(3)$ spin chain sigma model and string theory,” *JHEP* **0404**, 052 (2004) [arXiv:hep-th/0403139].
- [36] S. Bellucci, P. Y. Casteill, J. F. Morales and C. Sochichiu, “ $\text{SL}(2)$ spin chain and spinning strings on $\text{AdS}(5) \times S^{*5}$,” *Nucl. Phys. B* **707**, 303 (2005) [arXiv:hep-th/0409086].
- [37] S. A. Frolov, R. Roiban and A. A. Tseytlin, “Gauge - string duality for superconformal deformations of $N = 4$ super Yang-Mills theory,” *JHEP* **0507**, 045 (2005) [arXiv:hep-th/0503192].
- [38] S. Benvenuti and M. Kruczenski, “Semiclassical strings in Sasaki-Einstein manifolds and long operators in $N = 1$ gauge theories,” *JHEP* **0610**, 051 (2006) [arXiv:hep-th/0505046].
- [39] R. de Mello Koch, N. Ives, J. Smolic and M. Smolic, “Unstable giants,” *Phys. Rev. D* **73**, 064007 (2006) [arXiv:hep-th/0509007].
- [40] D. Berenstein, D. H. Correa and S. E. Vazquez, “A study of open strings ending on giant gravitons, spin chains and integrability,” [arXiv:hep-th/0604123].
- [41] R. de Mello Koch, J. Smolic and M. Smolic, “Giant gravitons - with strings attached. I,” arXiv:hep-th/0701066,
R. de Mello Koch, J. Smolic and M. Smolic, “Giant Gravitons - with Strings Attached (II),” *JHEP* **0709**, 049 (2007) [arXiv:hep-th/0701067],
D. Bekker, R. de Mello Koch and M. Stephanou, “Giant Gravitons - with Strings Attached (III),” *JHEP* **0802**, 029 (2008) [arXiv:0710.5372 [hep-th]].